

Character formulae for classical groups

P. E. Frenkel *

Abstract

We give formulae relating the value $\chi_\lambda(g)$ of an irreducible character of a classical group G to entries of powers of the matrix $g \in G$. This yields a far-reaching generalization of a result of J. L. Cisneros-Molina concerning the GL_2 case [C].

Introduction

The Weyl character formula tells us how to compute the character χ_λ of an irreducible finite dimensional representation V_λ with highest weight λ of a (complex, semisimple, connected) Lie group G :

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma z^{\sigma(\lambda+\rho)} = \chi_\lambda \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma z^{\sigma\rho}.$$

Here, for each weight $\ell \in \mathfrak{h}^*$ of the Lie algebra $\mathfrak{g}$ of G , the exponential $z^\ell : \tilde{H} \rightarrow \mathbb{C}^*$ is the corresponding multiplicative character of the preimage $\tilde{H}$ of a maximal torus $H \leq G$ in a universal covering $\tilde{G} \rightarrow G$. The weight $\rho \in \mathfrak{h}^*$ is the half-sum of the positive roots, $\mathfrak{W}$ is the Weyl group. Note that both sides of the formula are $\mathfrak{W}$ -antisymmetric characters of $\tilde{H}$, but χ_λ is well-defined as a $\mathfrak{W}$ -symmetric character of H . Note also that $\Delta = \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma z^{\sigma\rho}$ is not identically zero, so χ_λ is expressed as a ratio of Laurent polynomials in the coordinates on $\tilde{H}$. (We know *a priori* that χ_λ is itself a Laurent polynomial in the coordinates on H , but with many more terms in general than the numerator and denominator.)

The formula expresses the value of the character χ_λ at a group element $g \in G$ in terms of a conjugate of the semisimple part of g in the maximal torus H , i.e., in the case of the classical matrix groups, in terms of the eigenvalues of g . There are equally explicit expressions, called determinantal identities or Giambelli formulae [FH, Section A.1, formulae (A.5), (A.6) and Section A.3], in terms of the elementary resp. the complete symmetric polynomials in the eigenvalues.

In the present paper, we consider the connected classical groups and we prove variants of the Weyl character formula that express the value $\chi_\lambda(g)$ in many different, explicit rational ways in terms of the entries of powers of the generic matrix $g \in G$. It seems likely that these formulae provide the fastest and most straightforward way of calculating $\chi_\lambda(g)$ for generic g .

*Partially supported by OTKA grants T 042769 and T 046365.

This paper was motivated by J. L. Cisneros-Molina's paper [C] whose main result is the following. Let $\omega \neq 0$ be a linear function on the space M_2 of 2×2 matrices, such that $\omega(\mathbf{1}) = 0$. For example, ω could be one of the two off-diagonal entries. Then, for $\lambda = 0, 1, \dots$, we have $\omega(g^{\lambda+1})/\omega(g) = \text{tr } S^\lambda g$, the trace of the action of g on the λ -th symmetric power of the standard vector representation. Our results can be considered as far-reaching generalizations of this fact. In particular, for $g \in M_{r+1}$, and $\lambda = 0, 1, \dots$, the trace $\text{tr } S^\lambda g$ equals the ratio of the r -dimensional volumes of the two parallelepipeds spanned in M_{r+1}/M_1 by the images of $g, g^2, \dots, g^{r-1}, g^{\lambda+r}$ and of $g, g^2, \dots, g^{r-1}, g^r$, respectively (except when both volumes are zero, i.e., g has a minimal polynomial of degree $< r+1$). This is a particular case of Corollary 1.3 below.

Our proofs are motivated by the first of the four proofs given in [C], which is due to Jeremy Rickard.

1 General linear group

Let $G = GL_{r+1}(\mathbb{C})$ with $H = (\mathbb{C}^*)^{r+1}$ the maximal torus consisting of all invertible diagonal matrices, $\text{Hom}(H, \mathbb{C}^*) = \mathbb{Z}^{r+1}$ the weight lattice, and $\mathfrak{W} = \mathfrak{S}_{r+1}$ the Weyl group. Write

$$\rho = \left(\frac{r}{2}, \frac{r-2}{2}, \dots, \frac{2-r}{2}, \frac{-r}{2} \right)$$

for the half-sum of the positive roots. Set $\rho_t = \rho + (t, t, \dots, t, t)$ for $t \in \mathbb{C}$. For $\lambda = (\lambda_0, \dots, \lambda_r) \in \mathbb{Z}^{r+1}$, write $z^\lambda : H \rightarrow \mathbb{C}^*$ for the corresponding multiplicative character of the torus H , and, when λ is dominant, i.e. $\lambda_0 \geq \dots \geq \lambda_r$, write $\chi_\lambda : G \rightarrow \mathbb{C}$ for the character of the irreducible representation with highest weight λ . The Weyl character formula

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma z^{\sigma(\lambda + \rho_t)} = \chi_\lambda \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma z^{\sigma \rho_t}$$

holds with any $t \in \mathbb{C}$. The freedom in the choice of t comes from the central $\mathbb{C}^*$ in G . Both sides of the formula are $\mathfrak{W}$ -antisymmetric characters of the infinite cover $\tilde{H}$. When $\rho_t \in \mathbb{Z}^{r+1}$, both sides descend to H .

For $\ell \in \mathbb{C}^{r+1}$ and $g \in G$, define

$$g^\ell = \bigotimes_{i=0}^r g^{\ell_i} \in M_{r+1}(\mathbb{C})^{\otimes(r+1)}.$$

This is multi-valued, it depends on a choice of the value of $\log g \in \mathfrak{gl}_{r+1}(\mathbb{C})$. When $\ell \in \mathbb{Z}^{r+1}$, it is single-valued. When $\ell \in \mathbb{Z}_{\geq 0}^{r+1}$, we may allow $g \in M_{r+1}(\mathbb{C})$ rather than $g \in G$.

We have

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \bigwedge_{i=0}^r g^{\ell_i} \in M_{r+1}(\mathbb{C})^{\wedge(r+1)}.$$

Theorem 1.1 *Let $\lambda = (\lambda_0 \geq \dots \geq \lambda_r) \in \mathbb{Z}^{r+1}$ and $g \in GL_{r+1}(\mathbb{C})$. Then, for any $t \in \mathbb{C}$,*

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma(\lambda + \rho_t)} = \chi_\lambda(g) \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \rho_t};$$

equivalently,

$$\bigwedge_{i=0}^r g^{\lambda_i + r/2 - i + t} = \chi_\lambda(g) \cdot \bigwedge_{i=0}^r g^{r/2 - i + t},$$

where the powers are defined using any (but always the same) value of $\log g$. When $\rho_t \in \mathbb{Z}^{r+1}$, the powers are single-valued. In particular, for $t = r/2$, we get

$$\bigwedge_{i=0}^r g^{\lambda_i + r - i} = \chi_\lambda(g) \cdot \bigwedge_{i=0}^r g^{r - i}.$$

When ρ_t and λ are both in $\mathbb{Z}_{\geq 0}^{r+1}$, we may allow $g \in M_{r+1}(\mathbb{C})$.

Proof. The set of diagonalizable invertible matrices is dense in $M_{r+1}(\mathbb{C})$, so we may assume that g is such. The statement of the theorem is invariant under conjugation, so we may assume that $g = \text{diag}(z_0, \dots, z_r) \in H$. Then

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \bigwedge_{i=0}^r g^{\ell_i} = |z_j^{\ell_i}| \cdot \bigwedge_{j=0}^r e_{jj},$$

where e_{jj} is the diagonal matrix with a single 1 at the j -th position. The theorem now follows from the Weyl character formula. $\square$

Corollary 1.2 *Let Ω be an alternating $(r+1)$ -linear form on the space $M_{r+1}(\mathbb{C})$. Then, for $\lambda = (\lambda_0 \geq \dots \geq \lambda_r) \in \mathbb{Z}^{r+1}$, we have*

$$\Omega(g^{\lambda_0 + r}, g^{\lambda_1 + r - 1}, \dots, g^{\lambda_r}) = \chi_\lambda(g) \cdot \Omega(g^r, g^{r-1}, \dots, \mathbf{1}).$$

To express $\chi_\lambda(g)$ as a rational function in entries of powers of g , we must choose Ω such that the right hand side is not identically zero. For example, $\Omega(g_0, \dots, g_r)$ could be the determinant of the matrix formed by the diagonals, or by the first rows, etc. of the argument matrices. To calculate $\chi_\lambda(g)$ for a numerically given g , we need to choose Ω such that $\Omega(g^r, g^{r-1}, \dots, \mathbf{1}) \neq 0$. This is possible if and only if $\bigwedge_{i=0}^r g^{r-i} \neq 0$, i.e., the minimal polynomial of g is its characteristic polynomial. When g has a minimal polynomial of lower degree, we can use l'Hospital's rule.

Corollary 1.3 *Let ω be an alternating r -linear form on the space $M_{r+1}(\mathbb{C})$ such that ω vanishes if an argument is $\mathbf{1}$. Then, for λ as above and with $\lambda_r = 0$, we have*

$$\omega(g^{\lambda_0 + r}, g^{\lambda_1 + r - 1}, \dots, g^{\lambda_{r-1}}) = \chi_\lambda(g) \cdot \omega(g^r, g^{r-1}, \dots, g).$$

To express $\chi_\lambda(g)$ as a rational function in entries of powers of g , we must choose ω such that the right hand side is not identically zero. For example, $\omega(g_0, \dots, g_{r-1})$ could be the determinant of the $r \times r$ matrix formed by the truncated (i.e., leftmost entry omitted) first rows of the argument matrices. To calculate $\chi_\lambda(g)$ for a numerically given g , we need to choose ω such that $\omega(g^r, g^{r-1}, \dots, g) \neq 0$. This is possible if and only if the minimal polynomial of g is its characteristic polynomial.

Corollary 1.3, for $r = 1$, is the result of J. L. Cisneros-Molina's paper [C] mentioned in the Introduction.

To derive Corollary 1.3 from Corollary 1.2, simply set $\Omega = d\omega$, defined as usual by

$$\Omega(g_0, \dots, g_r) = \sum_{i=0}^r (-1)^i \omega(g_0, \dots, g_{i-1}, g_{i+1}, \dots, g_r).$$

Then $\Omega(g_0, \dots, g_{r-1}, \mathbf{1}) = (-1)^r \omega(g_0, \dots, g_{r-1})$ and Corollary 1.3 follows.

2 Special linear group

Let $G = SL_{r+1}(\mathbb{C})$ with $H \simeq (\mathbb{C}^*)^r$ the maximal torus consisting of all unimodular diagonal matrices, $\text{Hom}(H, \mathbb{C}^*) = \mathbb{Z}^{r+1}/\mathbb{Z}$ the weight lattice, and $\mathfrak{W} = \mathfrak{S}_{r+1}$ the Weyl group. Write $\rho = (r, r-1, \dots, 0) + \mathbb{Z} \cdot (1, \dots, 1) \in \mathbb{Z}^{r+1}/\mathbb{Z}$ for the half-sum of the positive roots. When $\lambda = (\lambda_0, \dots, \lambda_r) \in \mathbb{Z}^{r+1}/\mathbb{Z}$, write $z^\lambda : H \rightarrow \mathbb{C}^*$ for the corresponding multiplicative character of the torus H , and, when λ is dominant, write $\chi_\lambda : G \rightarrow \mathbb{C}$ for the character of the irreducible representation with highest weight λ . The Weyl character formula is valid as stated in the introduction. Both sides are $\mathfrak{W}$ -antisymmetric characters of H .

For $\ell \in \mathbb{Z}^{r+1}/\mathbb{Z}$ and $g \in G$, the antisymmetric tensor

$$\bigwedge_{i=0}^r g^{\ell_i} \in M_{r+1}(\mathbb{C})^{\wedge(r+1)}$$

is well defined because either g has a minimal polynomial of degree $< r+1$, in which case the algebra $\mathbb{C}[g]$ has dimension $< r+1$ and the antisymmetric tensor above is zero, or else g has its characteristic polynomial as minimal polynomial, in which case $\dim \mathbb{C}[g] = r+1$ and multiplication by g on it has determinant $\det g = 1$, so the tensor is independent of the chosen representative of ℓ .

Theorem 2.1 *Let $\lambda = (\lambda_0 \geq \dots \geq \lambda_r) + \mathbb{Z} \cdot (1, \dots, 1) \in \mathbb{Z}^{r+1}/\mathbb{Z}$ and $g \in SL_{r+1}(\mathbb{C})$. Then*

$$\bigwedge_{i=0}^r g^{\ell_i} = \chi_\lambda(g) \cdot \bigwedge_{i=0}^r g^{r-i},$$

where $\ell_i = \lambda_i + r - i$.

Proof. The theorem trivially follows from Theorem 1.1. □

3 Odd special orthogonal group

Let $G = SO_{2r+1}(\mathbb{C})$ be the connected group preserving the quadratic form

$$x_1 y_1 + \dots + x_r y_r + z^2.$$

We take the maximal torus $H = (\mathbb{C}^*)^r$ consisting of all special orthogonal diagonal matrices $\text{diag}(z_1, z_1^{-1}, \dots, z_r, z_r^{-1}, 1)$. In the weight lattice $\text{Hom}(H, \mathbb{C}^*) = \mathbb{Z}^r$, we take

$\lambda = (\lambda_1, \dots, \lambda_r)$ to correspond to the monomial $z^\lambda = \prod z_j^{\lambda_j}$. The Weyl group $\mathfrak{W}$ is the semidirect product of $\mathfrak{S}_r$ and Z_2^r . Write $\rho = (r - 1/2, r - 3/2, \dots, 3/2, 1/2)$ for the half-sum of the positive roots. The Weyl character formula is valid as stated in the introduction. Both sides are $\mathfrak{W}$ -antisymmetric characters of the double cover $\tilde{H}$.

For $\ell \in (\mathbb{Z} + \frac{1}{2})^r$ and $g \in G$, define

$$g^\ell = \bigotimes_{i=1}^r g^{\ell_i} \in M_{2r}(\mathbb{C})^{\otimes r}.$$

This is multi-valued, it depends on a choice of $\sqrt{g}$. We have

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) \in M_{2r}(\mathbb{C})^{\wedge r}.$$

Theorem 3.1 *Let $\lambda = (\lambda_1 \geq \dots \geq \lambda_r) \in \mathbb{Z}_{\geq 0}^r$ and $g \in SO_{2r+1}(\mathbb{C})$. Then*

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \chi_\lambda(g) \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \rho};$$

equivalently,

$$\bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) = \chi_\lambda(g) \cdot \bigwedge_{i=1}^r (g^{r+1/2-i} - g^{-(r+1/2-i)}),$$

where $\ell = \lambda + \rho$, i.e. $\ell_i = \lambda_i + r + 1/2 - i$, and the powers are defined using any, but always the same value of $\sqrt{g} \in SO_{2r+1}(\mathbb{C})$.

Proof. The set of diagonalizable matrices is dense in G , so we may assume that $\sqrt{g}$ is such. The statement of theorem is invariant under conjugation, so we may assume that

$$\sqrt{g} = \text{diag} \left(z_1^{1/2}, z_1^{-1/2}, \dots, z_r^{1/2}, z_r^{-1/2}, 1 \right) \in H.$$

Then

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) = |z_j^{\ell_i} - z_j^{-\ell_i}| \cdot \bigwedge_{j=1}^r (e_{jj} - f_{jj}),$$

where e_{jj} resp. f_{jj} is the diagonal matrix with a single 1 at the position corresponding to the x_j resp. y_j coordinate. The theorem now follows from the Weyl character formula. $\square$

4 Symplectic group

Let $G = Sp_{2r}(\mathbb{C})$ be the group preserving the skew bilinear form $\sum_{i=1}^r (x'_i y''_i - y'_i x''_i)$ on $\mathbb{C}^{2r}$. We take the maximal torus $H = (\mathbb{C}^*)^r$ consisting of all symplectic diagonal matrices $\text{diag}(z_1, z_1^{-1}, \dots, z_r, z_r^{-1})$. In the weight lattice $\text{Hom}(H, \mathbb{C}^*) = \mathbb{Z}^r$, we take $\lambda = (\lambda_1, \dots, \lambda_r)$ to correspond to the monomial $z^\lambda = \prod z_j^{\lambda_j}$. The Weyl group $\mathfrak{W}$ is the semidirect product of $\mathfrak{S}_r$ and Z_2^r . Write $\rho = (r, r - 1, \dots, 1)$ for the half-sum of the positive roots. The Weyl

character formula is valid as stated in the introduction. Both sides are $\mathfrak{W}$ -antisymmetric characters of H .

For $\ell \in \mathbb{Z}^r$ and $g \in G$, define

$$g^\ell = \bigotimes_{i=1}^r g^{\ell_i} \in M_{2r}(\mathbb{C})^{\otimes r}.$$

Then

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) \in M_{2r}(\mathbb{C})^{\wedge r}.$$

Theorem 4.1 *Let $\lambda = (\lambda_1 \geq \dots \geq \lambda_r) \in \mathbb{Z}_{\geq 0}^r$ and $g \in Sp_{2r}(\mathbb{C})$. Then*

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \chi_\lambda(g) \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \rho};$$

equivalently,

$$\bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) = \chi_\lambda(g) \cdot \bigwedge_{i=1}^r (g^{r+1-i} - g^{-(r+1-i)}),$$

where $\ell = \lambda + \rho$, i.e., $\ell_i = \lambda_i + r + 1 - i$.

Proof. The set of diagonalizable matrices is dense in G , so we may assume that g is such. The statement of theorem is invariant under conjugation, so we may assume that $g = \text{diag}(z_1, z_1^{-1}, \dots, z_r, z_r^{-1}) \in H$. Then

$$\sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} = \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) = |z_j^{\ell_i} - z_j^{-\ell_i}| \cdot \bigwedge_{j=1}^r (e_{jj} - f_{jj}),$$

where e_{jj} resp. f_{jj} is the diagonal matrix with a single 1 at the position corresponding to the x_j resp. y_j coordinate. The theorem now follows from the Weyl character formula. $\square$

5 Even special orthogonal group

Let $G = SO_{2r}(\mathbb{C})$ be the connected group preserving the quadratic form

$$Q = x_1 y_1 + \dots + x_r y_r.$$

We take the maximal torus $H = (\mathbb{C}^*)^r$ consisting of all special orthogonal diagonal matrices $\text{diag}(z_1, z_1^{-1}, \dots, z_r, z_r^{-1})$. In the weight lattice $\text{Hom}(H, \mathbb{C}^*) = \mathbb{Z}^r$, we take $\lambda = (\lambda_1, \dots, \lambda_r)$ to correspond to the monomial $z^\lambda = \prod z_j^{\lambda_j}$. The Weyl group $\mathfrak{W}$ is the semidirect product of $\mathfrak{S}_r$ and Z_2^{r-1} . It acts by permuting the indices and by performing an even number of sign changes. Write $\tilde{\mathfrak{W}} > \mathfrak{W}$ for the Weyl group in the full orthogonal group $O_{2r}(\mathbb{C})$. It is the semidirect product of $\mathfrak{S}_r$ and Z_2^r . If $\sigma \in \mathfrak{W}$, we write $[\sigma]$ for its image in $\mathfrak{S}_r$. Write $\rho = (r-1, r-2, \dots, 1, 0)$ for the half-sum of the positive roots. Write $\epsilon = (1, 1, \dots, 1, 1)$

so that $e = \epsilon + \rho = (r, r-1, \dots, 2, 1)$ is regular for $\tilde{\mathfrak{W}}$. The Weyl character formula is valid as stated in the introduction. Both sides are $\mathfrak{W}$ -antisymmetric characters of H .

For $\ell \in \mathbb{Z}^r$ and $g \in G$, define

$$g^\ell = \bigotimes_{i=1}^r g^{\ell_i} \in M_{2r}(\mathbb{C})^{\otimes r}.$$

We have

$$\begin{aligned} 2 \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \ell} &= \sum_{\sigma \in \tilde{\mathfrak{W}}} \left((-1)^{[\sigma]} + (-1)^\sigma \right) g^{\sigma \ell} = \\ &= \bigwedge_{i=1}^r (g^{\ell_i} + g^{-\ell_i}) + \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) \in M_{2r}(\mathbb{C})^{\wedge r}. \end{aligned}$$

Note that the second term is zero if any ℓ_i is zero.

Theorem 5.1 *Let $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathbb{Z}^r$ with $\lambda_1 \geq \dots \geq \lambda_{r-1} \geq |\lambda_r|$. Set $\bar{\lambda} = (\lambda_1, \dots, \lambda_{r-1}, -\lambda_r)$. Let $g \in SO_{2r}(\mathbb{C})$. Then*

$$2 \sum_{\sigma \in \tilde{\mathfrak{W}}} (-1)^{[\sigma]} g^{\sigma \ell} = (\chi_\lambda + \chi_{\bar{\lambda}})(g) \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^{[\sigma]} g^{\sigma \rho};$$

equivalently,

$$2 \bigwedge_{i=1}^r (g^{\ell_i} + g^{-\ell_i}) = (\chi_\lambda + \chi_{\bar{\lambda}})(g) \cdot \bigwedge_{i=1}^r (g^{r-i} + g^{-(r-i)}).$$

Also,

$$(\chi_\epsilon - \chi_{\bar{\epsilon}})(g) \cdot \sum_{\sigma \in \tilde{\mathfrak{W}}} (-1)^\sigma g^{\sigma \ell} = (\chi_\lambda - \chi_{\bar{\lambda}})(g) \cdot \sum_{\sigma \in \mathfrak{W}} (-1)^\sigma g^{\sigma \epsilon};$$

equivalently,

$$\sqrt{-1}^r \text{Pf}(g - g^{-1}) \cdot \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) = (\chi_\lambda - \chi_{\bar{\lambda}})(g) \cdot \bigwedge_{i=1}^r (g^{r+1-i} - g^{-(r+1-i)}).$$

Throughout, $\ell = \lambda + \rho$, i.e., $\ell_i = \lambda_i + r - i$.

Note that

$$(\chi_\epsilon - \chi_{\bar{\epsilon}})(g) = \sqrt{-1}^r \text{Pf}(g - g^{-1}),$$

where the sign of the Pfaffian of $g - g^{-1} \in \mathfrak{so}_{2r}(\mathbb{C})$ is specified by declaring the ordered Q -orthonormal bases of the standard vector representation $\mathbb{C}^{2r}$ with determinant $(2\sqrt{-1})^r$ to be of positive orientation.

Proof. The set of diagonalizable matrices is dense in G , so we may assume that g is such. The statement of the theorem is invariant under conjugation, so we may assume that $g = \text{diag}(z_1, z_1^{-1}, \dots, z_r, z_r^{-1}) \in H$. Then

$$\sum_{\sigma \in \tilde{\mathfrak{W}}} (-1)^{[\sigma]} g^{\sigma \ell} = \bigwedge_{i=1}^r (g^{\ell_i} + g^{-\ell_i}) = |z_j^{\ell_i} + z_j^{-\ell_i}| \cdot \bigwedge_{j=1}^r (e_{jj} + f_{jj})$$

and

$$\sum_{\sigma \in \tilde{\mathfrak{W}}} (-1)^{\sigma} g^{\sigma \ell} = \bigwedge_{i=1}^r (g^{\ell_i} - g^{-\ell_i}) = |z_j^{\ell_i} - z_j^{-\ell_i}| \cdot \bigwedge_{j=1}^r (e_{jj} - f_{jj}),$$

where e_{jj} resp. f_{jj} is the diagonal matrix with a single 1 at the position corresponding to the x_j resp. y_j coordinate. The theorem now follows from the Weyl character formula. $\square$

References

[C] J. L. Cisneros-Molina, An invariant of 2×2 matrices, *Electronic Journal of Linear Algebra* 13 (2005), 146–152.

[FH] W. Fulton and J. Harris, *Representation theory*, GTM, Springer, New York, 1991.